

Cube-magic labelings of grids

Rachel Wulan Nirmalasari Wijaya, Joe Ryan, and Thomas Kalinowski

University of Newcastle, Australia

10th February 2017

Abstract

We show that the vertices and edges of a d -dimensional grid graph $G = (V, E)$ ($d \geq 2$) can be labeled with the integers from $\{1, \dots, |V|\}$ and $\{1, \dots, |E|\}$, respectively, in such a way that for every subgraph H isomorphic to a d -cube the sum of all the labels of H is the same. As a consequence, for every $d \geq 2$, every d -dimensional grid graph is Q_d -supermagic where Q_d is the d -cube.

1 Introduction

The graphs considered in this paper are finite, undirected and simple. For a graph G , we denote its vertex set by $V(G)$ and its edge set by $E(G)$. A *graph labeling*, as introduced in [8], is an assignment of integers to the vertices or edges, or both, subject to certain conditions. Over the years, a large variety of different types of graph labelings have been studied, see [2] for an extensive survey.

For a graph H , we say that a graph G admits an H -covering if every edge of G belongs to at least one subgraph of G which is isomorphic to H . A graph $G = (V, E)$ which admits an H -covering is called H -magic if there exists a bijection $F : V \cup E \rightarrow \{1, 2, \dots, |V| + |E|\}$ and a constant $c = c(F)$, which we call the H -magic sum of F , such that

$$\sum_{v \in V(H')} F(v) + \sum_{e \in E(H')} F(e) = c$$

for every subgraph $H' \subseteq G$ with $H' \cong H$. If in addition $F(V) = \{1, \dots, |V|\}$ then we say that the graph G is H -supermagic. The case where H is a single edge was studied in [1], and the general concept for arbitrary graphs H was introduced in [3]. Since then H -magic and H -supermagic labelings have been studied for a variety of graphs H ([4, 5, 6, 7, 9]).

In this paper we show that for an integer $d \geq 2$, a d -dimensional grid graph G is Q_d -supermagic where Q_d denotes the d -cube. For $d = 2$, the 2-cube is the same as a 4-cycle C_4 and our result is a consequence of Theorem 1 in [4] which gives sufficient conditions for the cartesian product of a graph and a path to be C_4 -supermagic.

The structure of the paper is as follows. In Section 2 we fix some notation and state our main result. Section 3 contains the proof which is by induction on the dimension d , where the base case $d = 2$ is contained in Section 3.1 and the induction step in Section 3.2.

2 Notation and main result

For integers $k \leq \ell$ we denote the sets $\{1, 2, \dots, k\}$ and $\{k, k+1, \dots, \ell\}$ by $[k]$ and $[k, \ell]$, respectively. For integers $d \geq 2$ and $n_1 \geq n_2 \geq \dots \geq n_d \geq 2$, let $\text{GRID}(d_1, \dots, n_d)$ denote the $n_1 \times \dots \times n_d$ -grid graph, i.e., the cartesian product of d paths of lengths $n_1, \dots, n_d$. In other words, the vertex set of $G(d_1, \dots, n_d)$ is $V = [n_1] \times \dots \times [n_d]$ and edge set

$$E = \left\{ \{\mathbf{x}, \mathbf{y}\} : \mathbf{x}, \mathbf{y} \in V \text{ and } \sum_{i=1}^d |x_i - y_i| = 1 \right\}.$$

The graph $\text{GRID}(2, 2, \dots, 2)$ is called the d -cube and will be denoted by Q_d .

To simplify the presentation of our proof we will label the vertices and the edges separately and then combine the labelings to obtain the Q_d -supermagic labeling. A vertex labeling $f : V \rightarrow \{1, 2, \dots, |V|\}$ for a graph $G = (V, E)$ is called H -magic if there exists a constant $c = c(f)$, called the H -magic sum of f such that

$$\sum_{v \in V(H')} f(v) = c$$

for every subgraph $H' \subseteq G$ with $H' \cong H$. Similarly, an edge labeling $g : E \rightarrow \{1, 2, \dots, |E|\}$ for a graph $G = (V, E)$ is called H -magic if there exists a constant $c' = c'(g)$, called the H -magic sum of g such that

$$\sum_{e \in E(H')} g(e) = c'$$

for every subgraph $H' \subseteq G$ with $H' \cong H$. An H -magic vertex labeling f and an H -magic edge labeling g with H -magic sums $c = c(f)$ and $c' = c'(g)$ can be combined to obtain an H -supermagic labeling F with H -supermagic sum $c + |E(H)||V(G)|$ by setting $F(v) = f(v)$ for all $v \in V(G)$ and $F(e) = g(e) + |V(G)|$ for all $e \in E(G)$.

Theorem 1. *Let $d \geq 2$ and $n_1 \geq n_2 \geq \dots \geq n_d \geq 2$ be positive integers, and let $G = \text{GRID}(n_1, \dots, n_d)$. Then G admits a Q_d -magic vertex labeling f and a Q_d -magic edge labeling g .*

Based on the observation about combining H -magic vertex and edge labelings we obtain the following corollary.

Corollary 1. *Let $d \geq 2$ and $n_1 \geq n_2 \geq \dots \geq n_d \geq 2$ be positive integers. Then $\text{GRID}(n_1, \dots, n_d)$ is Q_d -supermagic.*

3 Proof of the main result

We proceed by induction on d . In Section 3.1 we treat the base case $d = 2$, and present explicit vertex and edge labelings for grid graphs $\text{GRID}(n_1, n_2)$. In Section 3.2 we assume $d \geq 3$, and we describe how labelings f and g for $\text{GRID}(n_1, \dots, n_d)$ can be constructed from the labelings $\tilde{f}$ and $\tilde{g}$ for $\text{GRID}(n_1, \dots, n_{d-1})$.

3.1 The base case $d = 2$

In order to describe the labeling in a compact way we use $[P]$ to denote the indicator function for a statement P , i.e.,

$$[P] = \begin{cases} 1 & \text{if } P \text{ is true,} \\ 0 & \text{if } P \text{ is false.} \end{cases}$$

We define a vertex labeling $f : [n_1] \times [n_2] \rightarrow [n_1 n_2]$ by

$$f(i, j) = \begin{cases} (i-1)n_2 + j & \text{if } i \text{ is odd and } j \text{ is odd,} \\ (i-1)n_2 + (n_2 + 1 - j) & \text{if } i \text{ is even and } j \text{ is even,} \\ (n_1 - i)n_2 + j + [2 \mid n_1 \wedge 2 \nmid n_2] & \text{if } i \text{ is odd and } j \text{ is even,} \\ (n_1 - i)n_2 + (n_2 + 1 - j + [2 \mid n_1 \wedge 2 \nmid n_2]) & \text{if } i \text{ is even and } j \text{ is odd,} \end{cases} \quad (1)$$

and an edge labeling $g : E \rightarrow [2n_1n_2 - n_1 - n_2]$ by

$$g((i, j), (i, j + 1)) = (i - 1)(2n_2 - 1) + j \quad \text{for } (i, j) \in [n_1] \times [n_2 - 1], \quad (2)$$

$$g((i, j), (i + 1, j)) = (n_1 - i)(2n_2 - 1) + 1 - j \quad \text{for } (i, j) \in [n_1 - 1] \times [n_2]. \quad (3)$$

Example 1. The construction is illustrated in Figure 1 for the graph $\text{GRID}(5, 3)$.




Figure 1: A C_4 -supermagic labeling for GRID(5,3). The vertex labels $f(i, j)$ are given by (1), and the edge labels are $g(e) + 15$ where g is defined by (2) and (3).

Lemma 1. *The function f defined by (1) is a Q_2 -magic vertex labeling for $\text{GRID}(n_1, n_2)$ with Q_2 -magic sum*

$$c(f) = \begin{cases} 2(n_1 n_2 + 1) & \text{if } n_1 \text{ is odd or } n_2 \text{ is even,} \\ 2(n_1 n_2 + 2) & \text{if } n_1 \text{ is even and } n_2 \text{ is odd.} \end{cases}$$

Proof. For every $(i, j) \in [n_1 - 1] \times [n_2 - 1]$ we have

$$\begin{aligned}
& f(i, j) + f(i, j + 1) + f(i + 1, j) + f(i + 1, j + 1) \\
&= (i - 1)n_2 + j + (n_1 - i)n_2 + (j + 1) + [2 \mid n_1 \wedge 2 \nmid n_2] + (n_1 - (i + 1))n_2 \\
&\quad + n_2 + 1 - (j + 1) + [2 \mid n_1 \wedge 2 \nmid n_2] + ((i + 1) - 1)n_2 + n_2 + 1 - (j + 1) \\
&= (2n_1 - 2)n_2 + 2n + 2 + 2[2 \mid n_1 \wedge 2 \nmid n_2] \\
&= 2(n_1n_2 + 1 + [2 \mid n_1 \wedge 2 \nmid n_2]).
\end{aligned}$$

Lemma 2. *The function g defined by (2) and (3) is a Q_2 -magic edge labeling for $\text{GRID}(n_1, n_2)$ with Q_2 -magic sum $c(g) = (2n_1 - 1)(2n_2 - 1) + 1$.*

Proof. For every $(i, j) \in [n_1 - 1] \times [n_2 - 1]$, we have

$$\begin{aligned} & g((i, j), (i, j + 1)) + g((i, j), (i + 1, j)) + g((i + 1, j), (i + 1, j + 1)) + g((i, j + 1), (i + 1, j + 1)) \\ &= (i - 1)(2n_2 - 1) + j + (n_1 - i)(2n_2 - 1) + 1 - j + ((i + 1) - 1)(2n_2 - 1) + j \\ &\quad + (n_1 - i)(2n_2 - 1) + 1 - (j + 1) \\ &= (2n_1 - 1)(2n_2 - 1) + 1. \end{aligned}$$

Combining these two labelings as described in Section 2 we obtain a Q_2 -supermagic labeling F with Q_2 -supermagic sum

$$c(F) = 10n_1n_2 - 2n_1 - 2n_2 + \begin{cases} 4 & \text{if } n_1 \text{ is odd or } n_2 \text{ is even,} \\ 6 & \text{if } n_1 \text{ is even and } n_2 \text{ is odd.} \end{cases}$$

3.2 The induction step

We now assume $d \geq 3$. By induction, there exist Q_{d-1} -magic labelings $\tilde{f}$ and $\tilde{g}$ with Q_{d-1} -magic sums $S = c(\tilde{f})$ and $S' = c'(\tilde{g})$ for $\text{GRID}(n_1, \dots, n_{d-1})$. We define $f : [n_1] \times \dots \times [n_d] \rightarrow \{1, \dots, \prod_{i=1}^d n_i\}$ by

$$f(x_1, \dots, x_d) = \begin{cases} \tilde{f}(x_1, \dots, x_{d-1}) + (x_d - 1) \prod_{i=1}^{d-1} n_i & \text{if } x_1 + \dots + x_{d-1} \text{ is even,} \\ \tilde{f}(x_1, \dots, x_{d-1}) + (n_d - x_d) \prod_{i=1}^{d-1} n_i & \text{if } x_1 + \dots + x_{d-1} \text{ is odd.} \end{cases} \quad (4)$$

Example 2. The vertex labeling (4) is illustrated in Figure 2 for the graph $\text{GRID}(5, 3, 3)$ where $\tilde{f}$ is the vertex labeling for $\text{GRID}(5, 3)$ presented in Example 1.




Figure 2: A Q_3 -magic vertex labeling with magic sum $c(f) = 184$ for $\text{GRID}(5, 3, 3)$.

Lemma 3. The function f defined by (4) is a Q_d -magic vertex labeling with Q_d -magic sum

$$c(f) = 2S + 2^{d-1}(n_d - 1) \prod_{i=1}^{d-1} n_i.$$

Before proving the lemma we illustrate the basic idea using the Q_3 -subgraph of $\text{GRID}(5, 3, 3)$ shown in Figure 3. This subgraph corresponds to the square labeled 12–7–8–5 in Figure 1, and




Figure 3: A vertex-labeled Q_3 -subgraph of $\text{GRID}(5, 3, 3)$. The triples next to the vertices indicate their position in the grid.

by (4) the vertex labels for the top and the bottom square of the cube in Figure 3 are

$$\begin{aligned} 27 &= 12 + 15, & 22 &= 7 + 15, & 23 &= 8 + 15, & 20 &= 5 + 15 \\ 42 &= 12 + 2 \times 15, & 7 &= 7, & 38 &= 8 + 2 \times 15, & 5 &= 5 \end{aligned}$$

and therefore the sum of the vertex labels is $2(12 + 7 + 8 + 5 + 4 \times 15) = 2S + 8 \times 15$, where S is the magic sum of the vertex labeling in Figure 1.

Proof of Lemma 3. Fix $i \in [n_d]$ and $(x_1, \dots, x_{d-1}) \in [n_1 - 1] \times \dots \times [n_{d-1} - 1]$. Let $H \cong Q_{d-1}$ be the subgraph of $\text{GRID}(n_1, \dots, n_d)$ induced by

$$V(H) = \left\{ (x_1 + \varepsilon_1, \dots, x_{d-1} + \varepsilon_{d-1}, i) : (\varepsilon_1, \dots, \varepsilon_{d-1}) \in \{0, 1\}^{d-1} \right\}.$$

Using the fact that exactly half of the 2^{d-1} vertices of H have even coordinate sum, we obtain from (4),

$$\begin{aligned} \sum_{v \in V(H)} f(v) &= \sum_{\varepsilon \in \{0, 1\}^{d-1}} \tilde{f}(x_1 + \varepsilon_1, \dots, x_{d-1} + \varepsilon_{d-1}) + 2^{d-2}(i-1) \prod_{i=1}^{d-1} n_i + 2^{d-2}(n_d - i) \prod_{i=1}^{d-1} n_i \\ &= S + 2^{d-2} \prod_{i=1}^{d-1} n_i (i-1 + n_d - i) = S + 2^{d-2}(n_d - 1) \prod_{i=1}^{d-1} n_i. \end{aligned} \quad (5)$$

A subgraph H of $\text{GRID}(n_1, \dots, n_d)$ is isomorphic to Q_d if and only if its vertex set is

$$V(H) = \{(x_1 + \varepsilon_1, \dots, x_{d-1} + \varepsilon_{d-1}, x_d + \varepsilon_d) : (\varepsilon_1, \dots, \varepsilon_{d-1}, \varepsilon_d) \in \{0, 1\}^d\}$$

for some $(x_1, \dots, x_d) \in [n_1 - 1] \times \dots \times [n_d - 1]$. Using (5), this implies

$$\sum_{v \in V(H)} f(v) = 2 \left(S + 2^{d-2}(n_d - 1) \prod_{i=1}^{d-1} n_i \right). \quad \square$$

We define an edge labeling $g : E \rightarrow \{1, \dots, |E|\}$ for $G = \text{GRID}(n_1, \dots, n_d)$ as follows. Let $N = |V(\text{GRID}(n_1, \dots, n_{d-1}))|$ and $M = |E(\text{GRID}(n_1, \dots, n_{d-1}))|$. For an edge $e = \{\mathbf{x}, \mathbf{y}\}$ with $y_d = x_d + 1$, we set

$$g(e) = \tilde{f}(x_1, \dots, x_{d-1}) + n_d M + \begin{cases} (x_d - 1)N & \text{if } x_1 + \dots + x_{d-1} \text{ is odd,} \\ (n_d - 1 - x_d)N & \text{if } x_1 + \dots + x_{d-1} \text{ is even.} \end{cases} \quad (6)$$

For the remaining edges we distinguish two cases.

Case 1. d is odd. For an edge $e = \{\mathbf{x}, \mathbf{y}\}$ with $y_i = x_i + 1$, $i \leq d - 1$, we set

$$g(e) = \tilde{g}(\{(x_1, \dots, x_{d-1}), (y_1, \dots, y_{d-1})\}) + \begin{cases} (x_d - 1)M & \text{if } i \text{ is odd,} \\ (n_d - x_d)M & \text{if } i \text{ is even.} \end{cases} \quad (7)$$

Case 2. d is even. For an edge $e = \{\mathbf{x}, \mathbf{y}\}$ with $y_i = x_i + 1$, $i \leq d - 2$, we set

$$g(e) = \tilde{g}(\{(x_1, \dots, x_{d-1}), (y_1, \dots, y_{d-1})\}) + \begin{cases} (x_d - 1)M & \text{if } i \text{ is odd,} \\ (n_d - x_d)M & \text{if } i \text{ is even.} \end{cases} \quad (8)$$

For an edge $e = \{\mathbf{x}, \mathbf{y}\}$ with $y_{d-1} = x_{d-1} + 1$, we set

$$g(e) = \tilde{g}(\{(x_1, \dots, x_{d-1}), (y_1, \dots, y_{d-1})\}) + \begin{cases} (x_d - 1)M & \text{if } \sum_{i=1}^{d-2} x_i \text{ is odd,} \\ (n_d - x_d)M & \text{if } \sum_{i=1}^{d-2} x_i \text{ is even.} \end{cases} \quad (9)$$

Example 3. The edge labeling given by (6) to (9) is illustrated in Figure 4 for the graph $\text{GRID}(5, 3, 3)$. The underlying labelings for $\tilde{f}$ and $\tilde{g}$ for $\text{GRID}(5, 3)$ are the labelings from Example 1.




Figure 4: A Q_3 -magic edge labeling with magic sum 594 for $\text{GRID}(5, 3, 3)$.

Lemma 4. The function g defined by (6) to (9) is a Q_d -magic edge labeling with Q_d -magic sum

$$c'(g) = S + 2S' + 2^{d-2}(n_d - 2)N + 2^{d-2}(2n_d + (d - 1)(n_d - 1))M.$$

Again, we illustrate the basic idea before going into the formal proof. Figure 5 shows the edge labels for the same subgraph of $\text{GRID}(5, 3, 3)$ as in the illustration for Lemma 3 (see Figure 3). The top and bottom squares of the cube in Figure 5 correspond to the square labeled 12–7–8–5 in Figure 6. In this example we have $N = 15$ and $M = 22$, and according to (6), the labels of the vertical edges are

$$78 = 12 + 3 \times 22, \quad 88 = 7 + 3 \times 22 + 15, \quad 74 = 8 + 3 \times 22, \quad 86 = 5 + 3 \times 22 + 15,$$





Figure 5: An edge-labeled Q_3 -subgraph of $\text{GRID}(5, 3, 3)$.

Figure 6: Q_2 -vertex magic and Q_2 -edge magic labeling of $\text{GRID}(5, 3)$.

so the sum of the labels of the vertical edges is $S + 12 \times 22 + 2 \times 15$, where $S = 12 + 7 + 8 + 5$ is the vertex-magic sum in Figure 6. According to (7), the labels of the edges in the top and the bottom square are

$$\begin{aligned} 15 &= 15, & 55 &= 11 + 2 \times 22, & 14 &= 14, & 50 &= 6 + 4 \times 22, \\ 37 &= 15 + 22, & 33 &= 11 + 22, & 36 &= 14 + 22, & 28 &= 6 + 22, \end{aligned}$$

and therefore the sum of these edge labels is $2(15 + 11 + 14 + 6 + 4 \times 22) = 2S' + 8 \times 22$, where $S' = 15 + 11 + 14 + 6$ is the edge-magic sum in Figure 6. Adding all edge labels we obtain $S + 2S' + 2 \times 15 + 20 \times 22$ as claimed in the lemma.

Proof of Lemma 4. Fix a subgraph $H \subseteq \text{GRID}(n_1, \dots, n_d)$ with $H \cong Q_d$. Its vertex set is

$$V(H) = \{(x_1 + \varepsilon_1, \dots, x_{d-1} + \varepsilon_{d-1}, x_d + \varepsilon_d) : (\varepsilon_1, \dots, \varepsilon_d) \in \{0, 1\}^d\}$$

for some $(x_1, \dots, x_d) \in [n_1 - 1] \times \dots \times [n_d - 1]$. We partition the vertex set as $V(H) = V_0(H) \cup V_1(H)$ and the edge set as $E(H) = E_0(H) \cup E_1(H) \cup E_2(H)$ where

$$\begin{aligned} V_\varepsilon(H) &= \{\mathbf{y} \in V(H) : y_d = x_d + \varepsilon\} & \text{for } \varepsilon \in \{0, 1\}, \\ E_\varepsilon(H) &= \{\{\mathbf{y}, \mathbf{z}\} \in E(H) : y_d = z_d = x_d + \varepsilon\} & \text{for } \varepsilon \in \{0, 1\}, \\ E_2(H) &= \{\{\mathbf{y}, \mathbf{z}\} \in E(H) : y_d = x_d, z_d = x_d + 1\}. \end{aligned}$$

Note that $|V_0(H)| = |V_1(H)| = |E_2(H)| = |V(Q_{d-1})| = 2^{d-1}$ and $|E_0(H)| = |E_1(H)| = |E(Q_{d-1})| = (d-1)2^{d-2}$. Using the fact that $y_1 + \dots + y_{d-1}$ is even for exactly half of the edges $\{\mathbf{y}, \mathbf{z}\} \in E_2(H)$ and that the subgraph of G induced by $V_0(H)$ is isomorphic to Q_{d-1} , we obtain from (6),

$$\begin{aligned} \sum_{e \in E_2(H)} g(e) &= \sum_{\mathbf{y} \in V_0(H)} \tilde{f}(y_1, \dots, y_{d-1}) + 2^{d-1}n_dM + 2^{d-2}(x_d - 1)N + 2^{d-2}(n_d - 1 - x_d)N \\ &= S + 2^{d-1}n_dM + 2^{d-2}(n_d - 2)N. \end{aligned} \quad (10)$$

For the edges in $E_0(H) \cup E_1(H)$ we use the fact that each of the sets $V_\varepsilon(H)$ induces a subgraph isomorphic to Q_{d-1} . In addition, if d is odd then exactly half of the indices $i \in \{1, \dots, d-1\}$ are even, and if d is even then

- exactly half of the indices $i \in \{1, \dots, d-2\}$ are even, and
- for exactly half of the vertices $(y_1, \dots, y_{d-2}, x_{d-1}, x_d + \varepsilon)$ the sum $y_1 + \dots + y_{d-2}$ is even.

In both cases we conclude that for exactly half of the edges in $E_\varepsilon(H)$ the term added to $\tilde{g}(\{(x_1, \dots, x_{d-1}), (y_1, \dots, y_{d-1})\})$ in (6) to (9) is $(x_d - 1)M$, and for the other half it is $(n_d - x_d)M$. This implies

$$\begin{aligned} \sum_{\{x,y\} \in E_\varepsilon(H)} g(\{x,y\}) &= \sum_{\{x,y\} \in E_\varepsilon(H)} \tilde{g}(\{(x_1, \dots, x_{d-1}), (y_1, \dots, y_{d-1})\}) \\ &\quad + (d-1)2^{d-3}(x_d - 1)M + (d-1)2^{d-3}(n_d - x_d)M \\ &= S' + (d-1)2^{d-3}(n_d - 1)M. \end{aligned} \quad (11)$$

Combining (10) and (11), the function g is a Q_d -magic labeling with Q_d -magic sum

$$\begin{aligned} c'(g) &= S + 2^{d-1}n_dM + 2 \left(S' + (d-1)2^{d-3}(n_d - 1)M \right) \\ &= S + 2S' + 2^{d-2}(n_d - 2)N + 2^{d-2}(2n_d + (d-1)(n_d - 1))M. \quad \square \end{aligned}$$

References

- [1] H. Enomoto, A. S. Lladó, T. Nakamigawa, and G. Ringel. Super edge-magic graphs. *SUT J. Math*, 34(2):105–109, 1998.
- [2] J. A. Gallian. A dynamic survey of graph labeling. *The electronic journal of combinatorics*, DS6(18th edition):1–389, 2015.
- [3] A. Gutiérrez and A. Lladó. Magic coverings. *Journal of Combinatorial Mathematics and Combinatorial Computing*, 55:43–56, 2005.
- [4] T. Kojima. On α -supermagic labelings of the cartesian product of paths and graphs. *Discrete Mathematics*, 313(2):164–173, 2013.
- [5] A. Lladó and J. Moragas. Cycle-magic graphs. *Discrete Mathematics*, 307(23):2925–2933, 2007.
- [6] T. Maryati, E. Baskoro, and A. Salman. P_h -supermagic labelings of some trees. *Journal of Combinatorial Mathematics and Combinatorial Computing*, 65:197–204, 2008.
- [7] A. Ngurah, A. Salman, and L. Susilowati. H -supermagic labelings of graphs. *Discrete Mathematics*, 310(8):1293–1300, 2010.
- [8] A. Rosa. On certain valuations of the vertices of a graph. In *Theory of Graphs (Internat. Symposium), Rome*, pages 349–355, 1966.
- [9] R. W. N. Wijaya, J. Ryan, and T. Kalinowski. H -supermagic labelings for firecrackers, banana trees and flowers. *arxiv:1607.07911*, 2016.